


# Detailed Ecological Risk Assessment: N.E. British Columbia: Wildlife Protection and Extractable Aliphatic Hydrocarbons

Craig Harris, M.Sc., P. Geo., R.P. Bio.

# Oil & Gas Commission – BC MoE Linkage



# Detailed Ecological Risk Assessment Content


---

- 1. Comprehensive Problem Formulation**
- 2. Comprehensive Understanding of Exposure**
- 3. Ecologically Relevant Effects Assessment**
- 4. Risk Characterization**
- 5. Uncertainty Analysis**

# Rationale for Selection of Wildlife Receptors

Receptor	Societal Value	Ecology/ Feeding Guild	Susceptibility/ Exposure	Habitat scale	Data Availability
Nelson's sharp-tail sparrow	BC Red Listed (high)	Migratory passerine insectivore	Ground nesting and ground foraging (high)	Small wetland areas	Limited
Caribou	BC Red Listed SARA Sch. 1 (high)	Ungulate mammalian herbivore	Observed feeding on site (high/moderate)	Large home range	Moderate/limited
Wolverine	BC Blue Listed Fur bearing (moderate)	Mammalian omnivore/carnivore	Contaminants not bioaccumulative (low)	Large home range	Limited
Meadow vole	(low)	Small mammalian herbivore/insectivore	Close contact with soil (high)	Small – within site	Good

# Conceptual Site Model



# Wildlife Exposure Models

---

$$ADD_{pot} = (\sum_{k=1}^m (C_k \times FS \times I_{df} \times FR_k)) / BW$$

Incidental soil

$$ADD_{pot} = (\sum_{k=1}^m (C_k \times I_{ff} \times FR_k)) / BW$$

Dietary food items

$$ADD_{adj} = SUF \times ADD_{pot}$$

Seasonal Adjusted Dose


# Do Aliphatic Hydrocarbons Transfer Through Food Chains ?

---

- MacLeod *et al.* (2004) \_ fugacity model predicting food chain transfer to fish, mammals and birds
  - No field validation and used carbon ranges too low ( $C_3 - C_{12}$ ) for weathered crude
- Chaineau *et al.* (1996) \_ field scale uptake study using 1.2% hydrocarbon contaminated soil
  - After 110 days of growth maize stems and leaves showed no detectable aliphatic or aromatic hydrocarbon
- Brandt *et al.* (2002) \_ field scale, multi year quantification of poly-aromatic hydrocarbons (PAH) in soil and terrestrial biota
  - mice and grasshoppers consistently displayed lower tissue PAH then frogs and vegetation

# Aliphatic Hydrocarbons in Food Chains: Salix leaves


ALS Sample ID: L1044695-7  
Client Sample ID: SALIX 11-07






# Aliphatic Hydrocarbons in Food Chains: Frogs

ALS Sample ID: L1044695-40  
Client Sample ID: ONSITE FROGS SMALL





# Aliphatic Hydrocarbons in Food Chains: Carex shoots

ALS Sample ID: L1044695-33  
Client Sample ID: CAREX 11-15



# Soil – Vegetation Tissue EPH Relationship



# Carex shoot EPH influenced by adsorbed peat ?

---



# Aliphatic Hydrocarbons: Wildlife Toxicity Reference Values (TRVs)

# EPH Wildlife Toxicity Reference Values

---

## **CCME CWS Basis for Livestock Protection:**

Stober (1962), Verträglichkeitsprüfungen mit roh und Heizöl an Rindern (*Research of cattle tolerability to raw and heating oil*)

## **CCME CWS Basis for Human Health Protection (aliphatic fractions c8-c16 & c17-c34)**

Edwards *et al*, (1997) , Total Petroleum Hydrocarbon Criteria Working Group Series, Volume 4



# Toxicological Studies Used in CCME TRVs

---

- Stober (1962) Stalled cattle orally dosed raw oil
  - Toxic threshold (weathered) = **8 mL / kg body weight**
  - Toxic threshold (fresh) = **2.5 mL / kg body weight**  $n = 1$
  - Toxic threshold (naptha) = **4 mL / kg body weight**
  - Effects on digestive system and liver
- Edwards *et al.* (1997) Rats orally dosed
  - Aliphatic c8-c17 hydrocarbon streams (unpublished)
    - Lowest observed effects = **500 mg / kg body weight**
    - Increased liver weight and size (reversible)
  - Aliphatic c17-c34 mineral oil (Smith *et al.* 1996)
    - Lowest observed effects = **2000 mg / kg body weight**
    - Mineral oil nodules in the liver tissue with associated inflammation


# Application of Uncertainty Factors (UF) in TRVs

All Lowest Observed Adverse Effects (LOAEL) data within the grey area.

Livestock TRV: Application of UF of 10 to Stober, 1962 LOAEL

Wildlife TRV: Apply range of UF to Edwards *et al*, 1997 LOAELs


Human TRV: Application of 100 – 1000 UF to Edwards *et al*, 1997 NOAELs






# Aliphatic Hydrocarbons and Wildlife Protection: Results

# Results: Incidental Soil Exposure



# Results: Incidental Soil + Dietary Exposure (vole)



Soil→biota EPH uptake factor of 0.14

Risk-based remediation currently under  
utilized in NE British Columbia

BC MoE process has well defined guidance and  
can be navigated by experienced professionals

Ecological risk decisions will continue to be driven by  
protection of eco-contact receptors and not wildlife

**Thank You**

Craig.Harris@aecom.com